

AD-A139 170

ELECTROCHEMICAL AND SPECTROSCOPIC STUDIES OF
POLYPYRIDINE COMPLEXES OF Fe^{+2} (U) YALE UNIV NEW HAVEN
CT DEPT OF EPIDEMIOLOGY AND PUBLIC HEALTH.
S SAHAMI ET AL. JAN 84 SUNYBUF/DC/TR-18


1/1

UNCLASSIFIED

F/G 7/4

NI





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER SUNYBUF/DC/TR-18	2. GOVT ACCESSION NO. AD-A239270	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Electrochemical and Spectroscopic Studies of Polypyridine Complexes of Fe(II)/(III) and Ru(II)/(III) in the Aluminum Chloride: N-(1-Butyl)Pyridinium Chloride Molten Salt	5. TYPE OF REPORT & PERIOD COVERED	
7. AUTHOR(s) Saeed Sahami and Robert A. Osteryoung	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	8. CONTRACT OR GRANT NUMBER(s) N00014-79-C-0682	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Arlington, VA 22117	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-715	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE January, 1984	
	13. NUMBER OF PAGES 39	
	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release: Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for Publication in Inorganic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Molten Salts; Voltammetry; Polypyridine Complexes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The electrochemical and spectroscopic behavior of $\text{Fe}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{phen})_3^{2+}$ and $\text{Ru}(\text{bpy})_3^{2+}$ in an ambient-temperature molten salt system, aluminum chloride:N-(1-butyl)pyridinium chloride (BuPyCl) have been studied as a function of melt composition. The complexes show a single one-electron reversible oxidation wave from melt compositions of 1:1 to 2:1 (AlCl_3 :BuPyCl mole ratio). Visible spect-		

DTIC

ELECTE

MAR 21 1984

A

AD A139170

DTIC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

roscopic and electrochemical data indicate that these compounds are stable in the neutral (1:1) and acid (mole ratio >1) melts, while $\text{Ru}(\text{bpy})_3^{2+}$ is stable in basic melts (mole ratio <1), but $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ decompose to FeCl_4^{2-} . Comparison of the formal potentials for M(III)/(II) -polypyridine couples studied here and in other solvents indicate that the redox properties of these complexes are relatively insensitive to the nature of the solvent. It was shown that oxygen acts as an oxidant in the acid melts to oxidize $\text{Fe}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{phen})_3^{2+}$ and $\text{Ru}(\text{bpy})_3^{2+}$ to their corresponding $3+$ form. The $3+$ form of each complex was stable in acid melts for several weeks. Perchlorate is also shown to function as an oxidant in acidic melts.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0682

Task No. NR-051-715

TECHNICAL REPORT NO. 18

ELECTROCHEMICAL AND SPECTROSCOPIC STUDIES OF POLYPYRIDINE COMPLEXES
OF Fe(II)/(III) and Ru(II)/(III) IN THE
ALUMINUM CHLORIDE: N-(1-Butyl)PYRIDINIUM CHLORIDE MOLTEN SALT

by

Saeed Sahami and Robert A. Osteryoung

Prepared for Publication in
Inorganic Chemistry

Department of Chemistry
State University of New York at Buffalo
Buffalo, New York 14214

January, 1984

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
AI	



Reproduction in whole or in part is permitted for any purpose of the
United States Government

Approved for Public Release: Distribution Unlimited

ABSTRACT

(2+)

(2+)

(2-)

The electrochemical and spectroscopic behavior of $\text{Fe}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{phen})_3^{2+}$ and $\text{Ru}(\text{bpy})_3^{2+}$ in an ambient-temperature molten salt system, aluminum chloride:N-(1-butyl)pyridinium chloride (BuPyCl) have been studied as a function of melt composition. The complexes show a single one-electron reversible oxidation wave from melt compositions of 1:1 to 2:1 (AlCl_3 :BuPyCl mole ratio). Visible spectroscopic and electrochemical data indicate that these compounds are stable in the neutral (1:1) and acid (mole ratio >1) melts, while $\text{Ru}(\text{bpy})_3^{2+}$ is stable in basic melts (mole ratio <1), but $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ decompose to FeCl_4^{2-} .

Comparison of the formal potentials for M(III)/(II)-polypyridine couples studied here and in other solvents indicate that the redox properties of these complexes are relatively insensitive to the nature of the solvent. It was shown that oxygen acts as an oxidant in the acid melts to oxidize $\text{Fe}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{phen})_3^{2+}$ and $\text{Ru}(\text{bpy})_3^{2+}$ to their corresponding 3+ form. The 3+ form of each complex was stable in acid melts for several weeks.

Perchlorate is also shown to function as an oxidant in acidic melts.

INTRODUCTION

The molten salt system composed of aluminum chloride and N-(1-butyl)pyridinium chloride (BuPyCl) is liquid at ambient temperatures ($\sim 30^\circ\text{C}$) over a wide compositional range varying from $\sim 0.7:1$ to $2:1$ (mole ratio of AlCl_3 to BuPyCl) (1,2). The Lewis acid-base properties of these melts change as the mole ratio of AlCl_3 to BuPyCl changes. The melts can be characterized as acidic, basic or neutral depending on the mole ratio of AlCl_3 to BuPyCl being greater than, less than or equal to unity (3). In the acidic melts, anionic species are Al_2Cl_7^- and AlCl_4^- , in basic melts AlCl_4^- and Cl^- , and in neutral, AlCl_4^- . It has been shown that equilibrium (1) with the $\log K = -16.9 \pm 2$ (4) provides



an adequate description of the system throughout the entire range of melt composition (2).

In view of their non-lability and extreme stability in the absence of direct illumination, low spin complexes of $\text{Ru}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ (where bpy = 2,2'-bipyridine and phen = 1,10-phenanthroline) were chosen to study in $\text{AlCl}_3\text{:BuPyCl}$ melts. These complexes were selected for several reasons. Initially we were interested to know whether these compounds are stable in this molten salt media, i.e. whether the reactant's coordination sphere remains intact while the Lewis acid-base properties of the melts are varied. A second aspect of this work was the examination of the possible use of these complexes as internal reference couples. In addition, the use of these melts as solvents for modified electrode studies, which includes Ru(II)/Ru(III) polymers, suggested such studies (5).

Spontaneous oxidation of hydrocarbons in highly acidic $\text{AlCl}_3\text{:BuPyCl}$ melts has been observed by Robinson and Osteryoung (1). The slow chemical oxidation of iodide in acidic $\text{AlCl}_3\text{:BuPyCl}$ melts has been suggested to be due to a reaction with traces of impurity present in the dry box atmosphere, probably molecular oxygen (6). To determine the true nature of the oxidizing species in the melt, these polypyridine complexes proved to be useful.

The electrochemistry of these transition-metal complexes has been studied both in protic and aprotic solvents (7-17). The oxidation states frequently observed for these complexes are 2+ and 3+. Polypyridine complexes of Ru and Fe with total charges of 1+, 0, and 1- have been observed in solvents with very negative potential windows such as acetonitrile (7,9) and N,N-dimethylformamide (8). Recently more highly oxidized species such as $\text{Ru}(\text{bpy})_3^{4+}$, $\text{Fe}(\text{bpy})_3^{4+}$ and $\text{Fe}(\text{bpy})_3^{5+}$ have been reported in liquid sulfur dioxide (13). We report here electrochemical and spectroscopic investigations of $\text{Fe}(\text{bpy})_3^{2+/3+}$, $\text{Fe}(\text{phen})_3^{2+/3+}$ and $\text{Ru}(\text{bpy})_3^{2+/3+}$ in ambient temperature $\text{AlCl}_3\text{:BuPyCl}$ ionic liquids.

EXPERIMENTAL SECTION

Preparation of N-(1-butyl)pyridinium chloride and purification of aluminum chloride have been described elsewhere (1). $\text{Ru}(\text{bpy})_3\text{Cl}_2 \cdot 6\text{H}_2\text{O}$, $\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$ and $\text{Fe}(\text{phen})_3(\text{ClO}_4)_2$ (G. F. Smith Chemical Co.), FeCl_2 (Alfa Products), FeCl_3 (Fisher), 2,2'-bipyridine (Baker) and 1,10-phenanthroline were used without further purification. Tetraethylammonium perchlorate (TEAP), (Baker) was dried in a vacuum oven and used for experiments testing the ability of perchlorate to function as an oxidant. Aluminum wire (Alfa Products) was cleaned in a 30:30:40 volume mixture of $\text{H}_2\text{SO}_4:\text{HNO}_3:\text{H}_3\text{PO}_4$, rinsed with water and dried. Chemicals were stored and all electrochemical experiments performed under argon atmosphere in a Vacuum Atmospheres Co. dry box.

A Metrohm glass cell covered with a teflon lid which had several holes for reference and counter electrode compartments, working electrode and thermometer was used for the electrochemical measurements. The entire cell assembly was placed in a furnace and the temperature controlled at $40 \pm 1^\circ\text{C}$ by a Thermo Electric Selector 800 temperature controller. Reference and counter electrode compartments were aluminum wires dipped into the 2:1 $\text{AlCl}_3:\text{BuPyCl}$ melt and both were separated from the working compartment by fine glass frits. A glassy carbon disc (GC) electrode obtained from Pine Instrument Co. was used as the working electrode. The GC (with the area of 0.196 cm^2 or 0.454 cm^2) was polished with successively finer grades of 1.0, 0.3 and 0.05 micron alumina (Buehler), then rinsed with water and air dried prior to transfer to the dry box.

All voltammograms were obtained using a EG&G PARC 175 Universal Programmer with a PARC 173 potentiostat and a Houston Omnigraph Model

2000 recorder. A Pine Instrument Co. electrode rotator (Model ASR2) was used for rotating disc electrode (RDE) studies.

UV and visible absorption spectra were recorded in either 0.1 or 1.0 cm path length quartz cells fitted with air-tight Teflon caps using a Perkin Elmer (Coleman 575) spectrophotometer. The cells were filled and sealed in the dry box.

RESULTS

1. ELECTROCHEMISTRY

a. OXIDATION OF $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$:

$\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$ and $\text{Fe}(\text{phen})_3(\text{ClO}_4)_2$ both dissolve in the neutral (mole ratio = 1.0) and acid (mole ratio >1) melts and give intense red-colored solutions. The solubility of $\text{Fe}(\text{phen})_3(\text{ClO}_4)_2$ is less than that of $\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$, especially in the neutral melt. In Figure 1, typical cyclic voltammograms at a GC electrode are shown for $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$, respectively. Cyclic and rotating disc electrode voltammograms for oxidation of polypyridine complexes of iron(II) were recorded as a function of melt acidity. Tabulations of cyclic and RDE voltammetric parameters, anodic to cathodic peak current ratios i_p^a/i_p^c , the difference between the anodic and cathodic peak potentials $\Delta E_p = E_p^a - E_p^c$, and $i_p^a/\nu^{1/2}$ for $\text{Fe}(\text{bpy})_3^{2+/3+}$ and $\text{Fe}(\text{phen})_3^{2+/3+}$ are presented in Tables 1 and 2, respectively. As shown in Figure 2 plots of the current vs. the square root of the electrode rotation rate at potentials on the rising portion of the waves and on the plateaus for oxidation of both complexes in the neutral melt were linear and passed through the origin, indicating that the oxidation is reversibly convective diffusion-controlled (18). Plots of E vs. $\log \frac{I_{\infty} - I}{I}$, constructed from RDE data for oxidation of both polypyridine complexes of Fe(II) were linear with slopes of 65 ± 1 mV ($2.3 \text{ RT/F} = 62$ mV at 40°C) (see Figure 3). Solutions of $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ in the neutral melt were found to be stable for a period of over a week. There was no change in the color of the solutions or their visible spectra. RDE voltammograms were recorded daily for polypyridine complexes

of Fe(II) in the neutral melt. The values of $E_{1/2}$ and anodic limiting current, I_1^a , were found to be constant within experimental error for at least seven days. When a neutral melt containing either of the polypyridine complexes of iron was made acidic by addition of $AlCl_3$, RDE voltammograms showed both cathodic and anodic current, indicating the presence of both the reduced and oxidized form of tris-bpy and tris-phen complexes of iron in the solution. Indeed in the 1.05:1 melt the cathodic limiting current, I_1^c , is <5% of the total limiting current, $I_1^t = I_1^c + I_1^a$. As the acidity of the melt increased the I_1^c/I_1^a ratio also increased and the color of the solution changed from red to dark brown and finally to blue. When the melt composition approached 1.9:1 almost all the complex was in the oxidized form. This oxidation of $Fe(bpy)_3^{2+}$ and $Fe(phen)_3^{2+}$ not only occurred with increased melt acidity (i.e., increase in $AlCl_3$ concentration), but as shown in Figure 4, even when a 1.2:1 melt containing 3.8 mM $Fe(bpy)_3^{2+}$ was kept in the dry box, I_1^c increased with time while I_1^a decreased. After about two hours I_1^c was ~ 3 times larger and I_1^a ~ 3 times smaller than their initial values. Thus, $\sim 75\%$ $Fe(bpy)_3^{2+}$ was oxidized to $Fe(bpy)_3^{3+}$ in less than two hours and eventually almost all of it was oxidized. It has been suggested that the slow chemical oxidation of I^- (with $E_{1/2} \approx 1$ V), in an acidic $AlCl_3$:BuPyCl melt, resulted from a reaction with traces of impurity present in the dry box atmosphere, possibly molecular oxygen (6). Although the dry box atmosphere had an oxygen level of ≤ 5 ppm, it may have been enough to slowly oxidize both $Fe(bpy)_3^{2+}$ and $Fe(phen)_3^{2+}$ complexes to their corresponding 3+ form, although perchlorate can also function as an oxidant (see below). The formation of the oxidized form of these complexes was proved both by RDE


experiments and their visible spectra (see below). Neither the reduction of oxygen nor electrochemistry from its reduction product could be seen directly by the electrochemical techniques. To show that O_2 was an oxidizing species, several other experiments were performed. Acidic $AlCl_3:BuPyCl$ melts with various compositions (1.07:1, 1.2:1 and 1.9:1) were prepared and $Fe(phen)_3(ClO_4)_2$ dissolved in them. These solutions were taken outside the dry box in sealed tubes. When these solutions were purged with oxygen, their color changed from red to blue (although the color change in the less acidic melt, 1.07:1 was slower than in the other melts). The spectra of these solutions were identical to those for acidic solutions of $Fe(phen)_3^{2+}$ which were oxidized slowly to $Fe(phen)_3^{3+}$ in the dry box. These results clearly indicate that oxygen in the acid melts can oxidize $Fe(bpy)_3^{2+}$ and $Fe(phen)_3^{2+}$ to $Fe(bpy)_3^{3+}$ and $Fe(phen)_3^{3+}$, respectively. The rate of the oxidation of these complexes appears to depend both on the amount of oxygen present in the dry box atmosphere and the melt acidity (see below, however). It is obvious from our results that the oxidation of any of these complexes in a given melt by traces of oxygen or perchlorate (see below) does not affect the E_f or $E_{1/2}$ values. The only changes were in the values of I_1^c and I_1^a . In any case the sum of $I_1^c + I_1^a$ (i.e., I_1^t), remained unchanged within experimental error. The oxidized forms of these polypyridine complexes of iron i.e., $Fe(bpy)_3^{3+}$ and $Fe(phen)_3^{3+}$ were stable in acid melts for at least two weeks.

The possibility that perchlorate could function as an oxidant in these systems was examined. As mentioned above, both $Fe(phen)_2^{2+}$ and

$\text{Fe}(\text{bpy})^{2+}$, were added to the melts as perchlorate salts and underwent slow oxidation in the acidic melts, some of which, as demonstrated above, was due to oxygen, but some of which might have been due to the presence of perchlorate. As described below, although the oxidation of the $\text{Ru}(\text{bpy})^{2+}$ complex was observed in very acidic (1.7:1) melts, it appeared much slower than that of the corresponding $\text{Fe}(\text{II})$ complexes. Since the $\text{Ru}(\text{bpy})^{2+}$ salts were chlorides, they were used in experiments to ascertain the effect of perchlorate as an oxidant. If a forty-fold excess of TEAP were added to a 1.7:1 $\text{AlCl}_3:\text{BuPyCl}$ melt containing ca 1 mM $\text{Ru}(\text{bpy})^{2+}$, the orange solution took about 30 minutes to change to green. This compares to much longer periods of time in the absence of perchlorate (see below). In a 1.1:1 melt, visible changes in color took place much more slowly; this was under conditions where, in the absence of perchlorate, no change in the solution color or electrochemistry was noted. In both basic and neutral (i.e., 1.0:1.0) melts, no indication of the oxidation of $\text{Ru}(\text{bpy})^{2+}$ was noted from spectroscopic measurements. These observations indicate that perchlorate, like oxygen, can act as an oxidant only in the acidic melt, that, qualitatively, the oxidation rate increases with increasing melt acidity and that there is no oxidation by perchlorate in the basic or neutral melt. Further experiments on this are in progress (19).

As is the case in other solvents, it was not possible to generate $\text{Fe}(\text{bpy})_3^{3+}$ or $\text{Fe}(\text{phen})_3^{3+}$ simply by adding ligands to an acidic solution of $\text{AlCl}_3:\text{BuPyCl}$ containing FeCl_3 . However, the polypyridine complexes of $\text{Fe}(\text{II})$ were easily obtained in acid melts by mixing FeCl_2 and ligands.

In basic $\text{AlCl}_3\text{:BuPyCl}$ melts (mole ratio <1), it was found that complexes of both $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ decompose and form FeCl_4^{2-} (20). Upon dissolving either of these complexes in the basic melt a red solution was obtained initially which changed within minutes to a light yellow color. Cyclic and RDE voltammograms for $\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$ in a 0.95:1 $\text{AlCl}_3\text{:BuPyCl}$ melt are shown in Figure 5. These voltammograms actually correspond to the oxidation of FeCl_4^{2-} to FeCl_4^- . There is excellent agreement between these results and those obtained by Nanjundiah et al., who studied the $\text{FeCl}_4^{2-}/1^-$ couple in basic $\text{AlCl}_3\text{:BuPyCl}$ melts (20). In the "neutral" melt, i.e., mole ratio = 1.0, where the concentration of Cl^- is negligible, both $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ are stable. We found, however, that the addition of even 20 mM BuPyCl (i.e., Cl^-) to a neutral melt containing a 3-4 mM of either of the polypyridine complexes of Fe(II) was sufficient to cause their dissociation to FeCl_4^{2-} . Although Cl^- is a relatively poor nucleophile in aqueous solutions, it seems to be highly reactive toward charged ions in aprotic solvents (21). An investigation of decomposition of tris-phenanthroline Fe(II) in dimethylsulfoxide (DMSO) showed that $\text{Fe}(\text{phen})_3^{2+}$ reacts rapidly with chloride (21,22). The reaction was assumed to be due to anation by chloride, leading to the formation of blue-colored $\text{Fe}(\text{phen})_2\text{Cl}_2$ complex (21). Thus, the complete dissociation of $\text{Fe}(\text{L})_3^{2+}$ (where L = bpy or phen) to FeCl_4^{2-} (see equation 2) in the basic $\text{AlCl}_3\text{:BuPyCl}$ is not surprising.



b. OXIDATION OF $\text{Ru}(\text{bpy})_3^{2+}$:

$\text{Ru}(\text{bpy})_3\text{Cl}_2 \cdot 6\text{H}_2\text{O}$ dissolves throughout the entire range of melt composition giving highly colored orange solutions. The electrochemical oxidation of this complex has been studied by cyclic and rotating disc voltammetry. A typical cyclic voltammogram at a GC electrode is shown in Figure 1. Cyclic and rotating disc electrode voltammograms for oxidation of $\text{Ru}(\text{bpy})_3^{2+}$ were examined as a function of melt composition from 0.8:1 to 2:1. Data obtained from these voltammograms are listed in Table 3. $\text{Ru}(\text{bpy})_3^{2+}$ showed no oxidation or reduction in the basic melt. In acidic or neutral melts only a single oxidation wave was obtained. Table 3 shows scan rate dependencies of peak potentials and peak currents for the $\text{Ru}(\text{bpy})_3^{2+/3+}$ couple. The ratio of one for i_p^a/i_p^c indicates that the electron transfer is reversible over the entire acidic melt composition. For a diffusion-controlled process, $i_p^a/\nu^{1/2}$ should be constant and independent of scan rate; this is true for the present case. Although ΔE_p increases slightly with the scan rate, it does not change with variation in the melt composition. The experimental peak separations (64-70 mV) correspond closely to the theoretical value for a one-electron process of 62 mV at 40°C. Plots of E vs. $\log [I_1 - I/I]$ in all melt compositions studied were linear with slopes of 65 ± 1 mV (see Figure 3).

As with the polypyridine complexes of Fe(II), RDE voltammograms showed that when the acidity of $\text{Ru}(\text{bpy})_3^{2+}$ containing melt increased there was a small cathodic current due to the reduction of $\text{Ru}(\text{bpy})_3^{3+}$

present in the solution. This cathodic current could only be observed when the melt acidity (mole ratio) was greater than $\sim 1.7:1$. Apparently the oxidation of $\text{Ru}(\text{bpy})_3^{2+}$ is slower than that of the polypyridine complexes of $\text{Fe}(\text{II})$. The initial I_1^c in a solution of $1.9:1 \text{ AlCl}_3:\text{BuPyCl}$ containing $\text{Ru}(\text{bpy})_3^{2+}$ was about 5% of the I_1^t . Even after two weeks the I_1^c was less than 55% of I_1^t , indicating that $\sim 55\%$ of $\text{Ru}(\text{bpy})_3^{2+}$ was oxidized to $\text{Ru}(\text{bpy})_3^{3+}$. On going from the reduced to the oxidized form of tris-bpy-ruthenium the color of the solution changed from orange to dusty brown and finally to green. As before the nature of the oxidant (i.e., traces of O_2 in the dry box atmosphere) was confirmed by purging acidic solutions of $\text{AlCl}_3:\text{BuPyCl}$ containing $\text{Ru}(\text{bpy})_3^{2+}$ with oxygen. Although it was possible to oxidize these solutions (in sealed tubes) over the melt compositions of $1.2:1$ to $1.9:1$ by oxygen, the time needed to convert $\text{Ru}(\text{bpy})_3^{2+}$ to $\text{Ru}(\text{bpy})_3^{3+}$ (i.e., change of color from orange to green) was longer as the acidity decreased. The green solutions of $\text{Ru}(\text{bpy})_3^{3+}$ was found to be stable for months.

c. AN ALUMINUM-OXYGEN CELL

In view of the observed oxidation of the polypyridine complexes of $\text{Fe}(\text{II})$ and $\text{Ru}(\text{II})$ by oxygen in acidic melts, an attempt was made to see if a cell using oxygen as an oxidant could be made. The reduction of O_2 by $\text{Ru}(\text{bpy})_3^{2+}$ is potentially a useful reaction for batteries since $\text{Ru}(\text{bpy})_3^{3+}$ is a strong and facile oxidizing agent. We have made a preliminary investigation (23) of this possibility by constructing the cell shown in Figure 6. Aluminum oxidation occurs at the anode and $\text{Ru}(\text{bpy})_3^{3+}$ reduction occurs at the cathode. The $\text{Ru}(\text{bpy})_3^{3+}$ is regenerated in the acid melt by oxygen and so $\text{Ru}(\text{bpy})_3^{2+}$ acts as a mediator for O_2 reduction at the

cathode. Thus the net cell reactions are Al oxidation and O_2 reduction; the product of O_2 reduction in the melt is not known at present, but is probably $AlOCl_2^-$. Reduction of $Ru(bpy)_3^{3+}$ by aluminum is extremely slow so no separator was necessary. The cell of Figure 6 gave an open circuit potential of ~ 1.4 V and passed a charge corresponding to 400 turnovers of the ruthenium complex with no significant deterioration. The potential is that which is expected based on the $Ru(bpy)_3^{+3/+2}$ voltammogram (see Figure 1C). The current was limited by mass transport of $Ru(bpy)_3^{3+}$ to the cathode.

2. SPECTROSCOPY

UV-visible absorption spectra for both the bivalent and the trivalent complexes of tris-bipyridine complexes of iron and ruthenium and tris-phenanthrolineiron in the neutral or acidic $AlCl_3:BuPyCl$ melts are shown in Figures 7-9. The trivalent compounds were obtained by the chemical oxidation of the respective divalent complexes in acid melts either by bubbling with oxygen or by trace of O_2 impurity present in the dry box atmosphere. The spectrum of $Ru(bpy)_3^{2+}$ did not change as the melt composition varied from basic to acidic. Thus $Ru(bpy)_3^{2+}$, unlike $Fe(bpy)_3^{2+}$ and $Fe(phen)_3^{2+}$ complexes, does not decompose in the basic melt. The spectra of $Fe(bpy)_3^{2+}$ and $Fe(phen)_3^{2+}$ in the neutral melt are identical to their respective spectrum in the slightly acidic melt. In Table 4 the molar absorptivities and maxima wavelengths obtained in this work are summarized and compared with those reported in other solvents (24-31). There is good agreement between positions of the absorption maxima determined here and those from the literature. The intense bands in the visible regions of these compounds are due to the transfer of electronic charge between the d-orbital of the metal ion and the

π -orbitals of the ligand. For the bivalent complexes charge transfer takes place from the metal to the ligand i.e., $t_{2g} \rightarrow \pi^*$ (MLCT) (24,25,30). In the case of trivalent complexes the charge transfer is from the ligand to the metal i.e., $\pi \rightarrow t_{2g}$ (LMCT) (27,31). In addition to these bands, there are other bands in the shorter wavelength portions of all spectra, which because of their higher intensities are outside the spectrum range. Due to the melt UV cutoff at ~ 280 nm, the UV absorption bands of these compounds could not be recorded.

DISCUSSION

Spectroscopic and electrochemical results show that all the complexes studied here are stable (i.e., have fixed coordination spheres) in the melt compositions of 1.0:1.0 to 2:1 $\text{AlCl}_3\text{:BuPyCl}$. Although $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ decompose in the basic melts to form FeCl_4^- , $\text{Ru}(\text{bpy})_3^{2+}$ is stable and does not dissociate, perhaps due to the significantly greater lability of the $\text{Fe}(\text{II})$ -tris complexes over that of ruthenium.

Based on the foregoing observations and results presented in Tables 1-3, we conclude that in the entire region from neutral to 2:1 melts the oxidation of $\text{Fe}(\text{bpy})_3^{2+}$, $\text{Fe}(\text{phen})_3^{2+}$ and $\text{Ru}(\text{bpy})_3^{2+}$ corresponds to a Nernstian one-electron electrochemically reversible process. Reversible behavior for these couples has been found in other solvents (8,12,13). Consideration of E_f or $E_{1/2}$ from Tables 1-3 indicates that although these values shift slightly toward more positive potentials with increasing melt acidity, the variation in E_f or $E_{1/2}$ in going from 1:1 to 1.9:1 mole ratio for all of these redox couples is less than 90 mV (the reproducibility of $\text{Al}/\text{Al}(\text{III})$ reference electrode is ± 10 mV). This indicates that the influence of variation in the melt composition on the redox properties of these polypyridine complexes is relatively small, but not as small as for ferrocene/ferricinium couple. Robinson and Osteryoung reported that the formal potential value of ferrocene/ferricinium in $\text{AlCl}_3\text{:BuPyCl}$ melts was independent of melt acidity (with the E_f value of 0.27 V vs. $\text{Al}/\text{Al}(\text{III})$) (1). This is understandable since the compact "sandwich" structure of ferrocene/ferricinium compared to the open structure of polypyridine complexes should prevent the close

approach of the solvent molecules to the metal redox center. A possible explanation of the observed variation in E_f for the polypyridine complexes with the melt acidity is the formation of ion-pairs, most probably with $AlCl_4^-$. As the acidity of the melt increases the concentration of $AlCl_4^-$ decreases and E_f becomes more positive. In other words with increasing acidity of the melt the extent of ion-pairing between the polypyridine complexes of iron and ruthenium with $AlCl_4^-$ decreases and the overall positive charge of the complex increases making it more difficult to oxidize.

Comparison of the E_f values for $Fe(bpy)_3^{2+/3+}$, $Fe(phen)_3^{2+/3+}$ and $Ru(bpy)_3^{2+/3+}$ (quoted vs. ferrocene/ferricinium) in different solvents with those in the neutral melt is shown in Table 5. It is obvious from this Table that indeed the redox properties of these couples are relatively insensitive to the nature of the solvent. This is expected since the polypyridine ligand should somewhat shield the solvent from the metal cation. Although the redox properties of these polypyridine couples appear to be largely independent of the solvent, indicating that they might be used as internal reference redox couples, they apparently do not offer any advantage over the ferrocene, particularly in $AlCl_3:BuPyCl$ melts.

In Table 6 values of diffusion coefficients for the polypyridine complexes of iron and ruthenium are given. The values of D were obtained from RDE experiments using the Levich equation:

$$I_l = 0.62 n F A D^{2/3} \omega^{1/2} \nu^{-1/6} C_0 \quad (3)$$

where I_l is the limiting current in amperes, n the number of electrons transferred, F the Faraday constant, A the area of the electrode in cm^2 ,


ω the rotation rate in radians/sec, ν the kinematic viscosity and C_0 the concentration of the electroactive species in mol/cm^3 (18). Diffusion coefficient for $\text{Fe}(\text{bpy})_3^{2+}$ and $\text{Fe}(\text{phen})_3^{2+}$ were only determined in the neutral melt where there was no chemical oxidation of the Fe(II) complex and its concentration was known accurately. The observed change in the D value for $\text{Ru}(\text{bpy})_3^{2+}$ in Table 6 can be attributed to changes in the viscosity (η) of the melt. The values of D and η are related by the modified Stokes-Einstein equation (33):

$$D = \frac{\kappa T}{4\pi\eta r} \quad (4)$$

where κ is the Boltzman constant and r is the radius of the diffusing species. The ηD values for $\text{Ru}(\text{bpy})_3^{2+}$ (tabulated in Table 6) are constant throughout the entire range of melt compositions studied here. Values of r calculated from equation (4) for polypyridine complexes are also listed in Table 6. These radii are in reasonable agreement with their corresponding crystallographic values (34,35), indicating that solvation is not very important in this medium for these compounds.

Although the "spontaneous" oxidation of some aromatic hydrocarbons (with $E_{1/2}$ of less than ~ 1.4 V) in highly acidic $\text{AlCl}_3:\text{BuPyCl}$ melts has been observed by Robinson and Osteryoung, the nature of the oxidizing species was not speculated upon (1). It is clear from this work that oxygen is an oxidant which oxidizes species with the $E_{1/2}$ of ~ 1.3 V and less in acidic $\text{AlCl}_3:\text{BuPyCl}$ melts. Considering our results and those observed previously for oxidations of aromatic hydrocarbons (1) and iodide (6), we conclude that O_2 can oxidize any species with an $E_{1/2}$ of 1.4 V or less in the acidic melts regardless of its chemical nature. In other words the oxidation process is insensitive to the nature of the

reductant and only depends on its formal potential, presumably according to the following reaction:



with $E_f \text{ O}_2/(\text{reduced O}_2) > E_f \text{ Red/Ox}$

Since neither the reduction of oxygen nor electrochemistry from its reduction product can be seen, it is not known whether the decrease in the rate of oxidation with decreasing melt acidity results from slow kinetics or unfavorable thermodynamics. One possible explanation is that by decreasing melt acidity, the value of E_f for $\text{O}_2/(\text{reduced O}_2)$ shifts to negative potentials. Indeed the absence of chemical oxidation of these compounds in the neutral or basic melts is a good indication of such an explanation. The other possibility is that the solubility of O_2 decreases drastically as the acidity decreases, making the oxidation more difficult.

It should also be pointed out that O_2 impurity in the dry-box may not be the sole trace oxidant present in these melts. Perchlorate, as indicated above, also functions as an oxidant in acidic melts. Recent work on the reduction of proton arising from the addition of water to the melt indicates that the potential for evolution of hydrogen moves in the positive direction with increasing melt acidity, suggesting that the "proton", which may arise from HCl or an $-\text{Al}-\text{O}-\text{H}$ entity formed as a result of water addition, is a stronger oxidant in acidic than basic melts (36). However, experiments at a platinum electrode in a melt containing both water and the Ru-complex indicated that the ruthenium reduction takes place at more positive potentials than does hydrogen evolution. Nevertheless, the ability of proton to act as an oxidant in these melts at the trace levels should not be overlooked.

ACKNOWLEDGEMENTS

We would like to thank Dr. Peter G. Pickup for helpful discussions and assistance. Helpful suggestions by other members of the molten salt research group, particularly Mr. T. Zawodzinski, Dr's. Z. Karpinski and M. Lipsztajn, are appreciated.

CREDIT

This work was supported by the Air Force Office of Scientific Research and the Office of Naval Research.

REFERENCES

1. Robinson, J.; Osteryoung, R. A., J. Am. Chem. Soc. 1979, 101, 323.
2. Gale, R. J.; Osteryoung, R. A., Inorg. Chem. 1979, 18, 1603.
3. Lipsztajn, M.; Osteryoung, R. A., J. Electrochem. Soc., 1983, 130, 1968.
4. Karpinski, Z. J.; Osteryoung, R. A., Inorg. Chem., in press.
5. Pickup, P. G.; Osteryoung, R. A., J. Electrochem. Soc. 1983, 130, 1965.
6. Karpinski, Z. J.; Osteryoung, R. A., J. Electroanal. Chem., in press.
7. Tokel, N. E.; Hemingway, R. E.; Bard, A. J., J. Am. Chem. Soc. 1973, 95, 6582.
8. Saji, T.; Aoyagui, S., J. Electroanal. Chem. 1975, 58, 401.
9. Saji, T.; Aoyagui, S., J. Electroanal. Chem. 1975, 60, 1.
10. Saji, T.; Aoyagui, S., J. Electroanal. Chem. 1975, 63, 31.
11. Tanaka, N.; Sato, Y., Electrochim. Acta 1968, 13, 335.
12. Mayer, U.; Kotocova, A.; Gutmann, V., J. Electroanal. Chem. 1979, 103, 409.
13. Gaudiello, J. G.; Sharp, P. R.; Bard, A. J., J. Am. Chem. Soc. 1982, 104, 6373.
14. Ogura, K.; Miyamoto, K., Electrochim. Acta 1978, 23, 509.
15. Samec, Z.; Nemec, I., J. Electroanal. Chem. 1971, 31, 161.
16. Sahami, S.; Weaver, M. J., J. Electroanal. Chem. 1981, 122, 155.
17. (a) Yee, E. L.; Cave, R. J.; Guyer, K. L.; Tyma, P. D.; Weaver, M. J., J. Am. Chem. Soc. 1979, 101, 1131.
(b) Yee, E. L.; Weaver, M. J., Inorg. Chem. 1980, 19, 1077.
18. Bard, A. J.; Faulkner, L. R., "Electrochemical Methods", 1980, John Wiley & Sons, Inc., New York, p. 288.
19. Sahami, S.; Osteryoung, R. A., unpublished results.
20. Nanjundiah, C.; Shimizu, K.; Osteryoung, R. A., J. Electrochem. Soc. 1982, 129, 2474.
21. Gillard, R. D.; Inorg. Chim. Acta 1979, 37, 103.

22. Farrington, D. J.; Jones, J. G.; Twigg, M. V., *Inorg. Chim. Acta* 1977, 25, L75.
23. Pickup, P. G.; Sahami, S.; Osteryoung, R. A., unpublished work.
24. Braddock, J. N.; Meyer, T. J., *J. Am. Chem. Soc.* 1973, 95, 3158.
25. Staniewicz, R. J.; Sympson, R. F.; Hendricker, D. G., *Inorg. Chem.* 1977, 16, 2166.
26. Bryant, G. M.; Fergusson, J. E.; Powell, H. K. J., *Aust. J. Chem.* 1971, 24, 257.
27. Wong, C. L.; Kochi, J. K., *J. Am. Chem. Soc.* 1979, 101, 5593.
28. Bryant, G. M.; Fergusson, J. E., *Aust. J. Chem.* 1971, 24, 275.
29. Nord, G.; Pederson, B.; Bjergbakke, E., *J. Am. Chem. Soc.* 1983, 105, 1913.
30. Schilt, A. A., "Analytical Applications of 1,10-Phenanthroline and Related Compounds", 1969, Pergamon Press, London, New York, p. 36.
31. Burgess, J.; Haines, R. J., *J. Inorg. Nucl. Chem.* 1977, 39, 1705.
32. McCaffery, A. J.; Mason, S. F.; Norman, B. J., *J. Chem. Soc. A.* 1969, 1428.
33. McLaughlin, E., *Trans. Faraday Soc.*, 1958, 55, 28.
34. Rillema, D. P.; Jones, D. S., *J. C. S. Chem. Comm.* 1979, 849.
35. Zalkin, A.; Templeton, D. H.; Ueki, T., *Inorg. Chem.* 1973, 12, 1641.
36. Sahami, S.; Osteryoung, R. A., *Anal. Chem.* 1983, 55, 1970.
37. Rillema, D. P.; Allen, G.; Meyer, T. J.; Conrad, D., *Inorg. Chem.* 1983, 22, 1617.

Table 1 Summary of Voltammetric Parameters for $\text{Fe}(\text{bpy})_3^{2+/3+}$ in AlCl_3 : BuPyCl Melts.^a

Melt Composition	Scan Rate(v) mV/s	Cyclic Voltammetric Data					RDE Data		
		E_p^c mV	E_a^d mV	E_f mV	ΔE_p mV	i_p^a/i_p^c	$i_p^a/\sqrt{v}^{1/2}$ mA cm ⁻² v ^{-1/2} s ^{1/2}	Rotation Rate rpm	$E_{1/2}$ mV
1:1	10	927	992	960	65	1.00	0.507	600	957
	20	927	992	960	65	1.01	0.510	900	959
	50	922	996	959	74	1.01	0.514	1600	960
	100	920	998	959	78	1.00	0.515	2000	959
								2500	963
1.05:1	10	935	1000	968	65	1.01	0.413	400	965
	20	935	1000	968	65	1.02	0.420	600	965
	50	936	1002	969	66	1.01	0.428	900	967
	100	931	1005	968	74	1.00	0.430	1600	970
	200	930	1010	970	80	1.00	0.433	2000	972
1.2:1	10	950	1015	982	65	1.01	0.451	600	980
	20	950	1015	982	65	1.01	0.478		
	50	948	1015	981	67	1.00	0.480		
	100	945	1020	982	75	1.01	0.494		
1.4:1	10	963	1025	994	62	1.01	0.407	600	990
	20	963	1025	994	62	1.00	0.412		
	50	960	1028	994	68	0.99	0.414		
	100	960	1030	995	70	1.02	0.414		
1.7:1	10	990	1055	1022	65	1.02	0.385	600	1020
	20	990	1055	1022	65	1.04	0.389		
	50	990	1057	1023	67	1.01	0.379		
	100	985	1060	1022	75	1.00	0.383		
1.9:1	10	1007	1073	1040	66	0.98	0.363	600	1038
	20	1007	1073	1040	66	0.99	0.358		
	50	1007	1075	1041	68	1.00	0.364		
	100	1005	1075	1040	70	0.99	0.369		

a. All potentials are in mV vs. $\text{Al}/\text{Al}(\text{III})$ in 2:1 AlCl_3 : BuPyCl, at $T=40^\circ\text{C}$.

b. Mole ratio of AlCl_3 : BuPyCl.

Table 2 Summary of Voltammetric Parameters for Fe(phen)₃^{2+/3+} in AlCl₃: BuPyCl Melts.^a

Melt Composition	Scan Rate (°) mV/s	Cyclic Voltammetric Data					RDE Data		
		E _p ^c mV	E _a ^p mV	E _f mV	ΔE _p mV	i _p ^a /i _p ^c	i _a ^a /i _p ^c 1/2	Rotation Rate rpm	E _{1/2} mV
1:1	10	965	1033	999	68	1.00	0.242	600	1002
	20	965	1035	1000	70	1.02	0.230	900	1005
	50	965	1035	1000	70	1.01	0.226	1600	1007
	100	962	1040	1001	78	1.01	0.233	2000	1005
1.05:1	10	976	1040	1008	64	1.02	0.268	600	1008
	20	975	1042	1008	67	1.00	0.288	900	1008
	50	973	1045	1009	72	1.01	0.293	1600	1007
	100	973	1047	1010	74	1.01	0.293	2500	1007
1.2:1	10	990	1052	1021	62	1.03	0.341	600	1020
	20	990	1055	1022	65	1.02	0.343	900	1021
	50	990	1058	1024	68	1.01	0.335	1600	1022
	100	988	1060	1024	72	1.00	0.338	2000	1025
1.4:1	10	1007	1070	1038	63	1.02	0.319	600	1038
	20	1007	1070	1038	63	1.02	0.323	900	1040
	50	1005	1072	1038	67	1.01	0.320	1600	1038
	100	1000	1075	1037	75	1.00	0.314	2000	1040
1.7:1	10	1035	1100	1067	65	1.01	0.270	600	1065
	20	1035	1100	1067	65	1.01	0.276	900	1065
	50	1032	1102	1067	70	1.00	0.280	1600	1065
	100	1032	1105	1068	73	1.00	0.275	2000	1065
1.9:1	10	1052	1118	1085	66	1.00	0.264	600	1083
	20	1052	1120	1086	68	1.01	0.261	900	1083
	50	1050	1120	1085	70	0.98	0.261	1600	1083
	100	1050	1120	1085	70	0.98	0.260	2000	1083

a. All potentials are in mV vs. Al/Al(III) in 2:1 AlCl₃: BuPyCl, at T=40°C.

b. Mole ratio of AlCl₃: BuPyCl

Table 3 Summary of Voltammetric Parameters for Ru(bpy)₃^{2+/3+} in AlCl₃: BuPyCl Melts.^a

Melt Composition	b	Scan Rate(ν) mV/s	Cyclic Voltammetric Data						RDE Data	
			E_p^c mV	E_a^d mV	E_f mV	ΔE_p^e mV	i_p^a/i_p^c	$i_p^a/\nu^{1/2}$ mA cm ⁻² V ^{-1/2} s ^{1/2}	Rotation Rate rpm	$E_{1/2}$ mV
1:1		10	1130	1198	1164	68	1.00	0.308	600	1163
		20	1130	1198	1164	68	1.03	0.300	900	1163
		50	1127	1202	1165	75	1.02	0.329	1600	1163
		100	1127	1204	1165	77	1.01	0.327	2000	1163
1.05:1		10	1140	1204	1172	64	1.00	0.357	400	1170
		20	1140	1204	1172	64	1.02	0.360	600	1170
		50	1138	1206	1172	68	1.03	0.364	900	1170
		100	1138	1208	1173	70	1.03	0.359	1600	1170
1.2:1		200	1136	1210	1173	74	1.03	0.353	2000	1170
									2500	1172
									3600	1175
									600	1183
1.4:1		10	1152	1216	1184	64	1.04	0.331		
		20	1152	1216	1184	64	1.06	0.329		
		50	1150	1218	1184	68	1.05	0.322		
		100	1150	1220	1185	70	1.05	0.321		
1.7:1		10	1170	1234	1202	64	1.02	0.344	600	1202
		20	1170	1234	1202	64	1.03	0.352		
		50	1170	1236	1204	68	1.03	0.343		
		100	1168	1240	1204	72	1.04	0.339		
1.9:1		10	1198	1262	1230	64	1.02	0.332	600	1230
		20	1198	1262	1230	64	1.03	0.338		
		50	1195	1265	1230	70	1.02	0.339		
		100	1195	1265	1230	70	1.04	0.340		
1.9:1		10	1220	1284	1252	64	1.02	0.331	600	1255
		20	1220	1284	1252	64	1.01	0.333		
		50	1220	1288	1254	68	1.02	0.325		
		100	1222	1290	1256	68	1.03	0.327		

a. All potentials are in mV vs. Al/Al(III) in 2:1 AlCl₃: BuPyCl, at T= 40° C

b. Mole ratio of AlCl₃: BuPyCl

Table 4. Spectroscopic Data for Polypyridine Complexes Of Ru and Fe in
 $\text{AlCl}_3\text{:BuPyCl}$ Melts.^a

Complex	color	λ_{max} , nm	molar abs. $\text{M}^{-1} \text{cm}^{-1}$	Values from the Literature
				λ_{max} , nm (molar abs., $\text{M}^{-1} \text{cm}^{-1}$)
$\text{Ru}(\text{bpy})_3^{2+}$	orange	454	14,000	453(13800) ^c , 450(14000) ^d
$\text{Ru}(\text{bpy})_3^{3+}$	green	424	3200	418(~3000) ^c
$\text{Fe}(\text{bpy})_3^{2+}$	red	523	8700	522(8740) ^e , 520(8240) ^f
$\text{Fe}(\text{bpy})_3^{3+}$	greenish-blue	627	380	618(284) ^g , 613(289) ^h
$\text{Fe}(\text{phen})_3^{2+}$	red	512	10,800	510(11300) ⁱ , 510(11100) ^j
$\text{Fe}(\text{phen})_3^{3+}$	blue	~610	800	590(600) ^j , 602(870) ^k

a. these spectra were obtained either in the neutral or acidic melts (see text for details)
 b. molar absorptivities are accurate to $\pm 5\%$.

- c. reference 24
- d. reference 25
- e. reference 26
- f. reference 27
- g. reference 13
- h. reference 28
- i. reference 29
- j. reference 30
- k. reference 31

Table 5. Formal Potentials for Polypyridine Complexes of Ru(II)/(III) and Fe(II)/(III) in Various Solvents vs. Ferrocene/Ferricinium Couple.

Solvent	E_f^a		
	$\text{Fe(phen)}_3^{2+/3+}$	$\text{Fe(bpy)}_3^{2+/3+}$	$\text{Ru(bpy)}_3^{2+/3+}$
Water	743 ^b	718 ^b	921 ^b
Acetonitrile	686 ^c	670 ^d	900 ^e
1:1 AlCl ₃ :BuPyCl	730 ^f	690 ^f	894 ^f
Nitromethane	721 ^c	727 ^d	

a. Formal potential for each redox couple is quoted in mV vs. E_f for Ferrocene/Ferricinium couple in same solvent i.e., $E_f = E_f(\text{redox couple}) - E_f(\text{Ferrocene/Ferricinium})$

b. Reference 17

c. Reference 12

d. Reference 16

e. Reference 37

f. This work

Table 6. Diffusion Coefficients (at $T = 40^\circ \text{C}$) and Radii of Polypyridine
Fe(II) and Ru(II) Complexes in $\text{AlCl}_3\text{:BuPyCl}$ Melts.^a

Melt Composition	$\text{Ru}(\text{bpy})_3^{2+}$		$\text{Fe}(\text{bpy})_3^{2+}$	$\text{Fe}(\text{phen})_3^{2+}$
	$D \times 10^7$ cm^2/s	$D \eta 10^8$ $\text{g cm}^2/\text{s}^2$	$D \times 10^7$ cm^2/s	$D \times 10^7$ cm^2/s
1:1	2.75	6.05	2.61	2.43
1.05:1	2.88	6.06		
1.2:1	2.99	6.00		
1.4:1	3.35	6.04		
1.7:1	3.67	6.05		
1.9:1	4.07	6.11		
Average = 6.04 0.04				
calculated radius of the complex, nm	0.57		0.60	0.64

a. Values of the viscosity (η) and density used in the calculation of D , $D\eta$ and radius are taken from reference 20.

FIGURE CAPTIONS

- Figure 1. Cyclic voltammograms in 1.05:1 $\text{AlCl}_3\text{:BuPyCl}$ melts. Scan rate = 50 mV/s; $T = 40^\circ\text{C}$. (a) 3.2 mM $\text{Fe}(\text{bpy})_3^{2+}$, GC electrode with $A = 0.454 \text{ cm}^2$. (b) 1.8 mM $\text{Fe}(\text{phen})_3^{2+}$, GC electrode with $A = 0.454 \text{ cm}^2$. (c) 2.8 mM $\text{Ru}(\text{bpy})_3^{2+}$, GC with $A = 0.196 \text{ cm}^2$.
- Figure 2. Plots of current at various potentials vs. $\omega^{1/2}$ for the oxidation of $\text{Fe}(\text{bpy})_3^{2+}$ (dashed lines), and $\text{Fe}(\text{phen})_3^{2+}$ (solid lines) in 1.0:1.0 $\text{AlCl}_3\text{:BuPyCl}$. E in mV: (1) 1000, (2) 1050, (3) 1300.
- Figure 3. Plots of E vs. $\log \frac{I_1 - I}{I}$ for the oxidation of (1) $\text{Fe}(\text{bpy})_3^{2+}$, (2) $\text{Fe}(\text{phen})_3^{2+}$, (3) $\text{Ru}(\text{bpy})_3^{2+}$ in 1.05:1 $\text{AlCl}_3\text{:BuPyCl}$.
- Figure 4. RDE voltammograms for a 1.2:1 $\text{AlCl}_3\text{:BuPyCl}$ melt with initial concentration of 3.8 mM $\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$, recorded as a function of time. Scan rate = 5 mV/s; rotation rate = 600 rpm; $T = 40^\circ\text{C}$; GC electrode with $A = 0.454 \text{ cm}^2$. Time in minutes: (1) 0, (2) 35, (3) 65, (4) 110, (5) 300.
- Figure 5. Voltammograms for 3.1 mM $\text{Fe}(\text{bpy})_3(\text{ClO}_4)_2$ dissolved in a 0.95:1 $\text{AlCl}_3\text{:BuPyCl}$ melt (which actually corresponds to 3.1 mM FeCl_4^{2-} , see text). $T = 40^\circ\text{C}$; GC electrode with $A = 0.454 \text{ cm}^2$. (a) Cyclic voltammogram with scan rate = 100 mV/s. (b) RDE voltammogram with scan rate = 5 mV/s and rotation rate = 600 rpm.
- Figure 6. An Aluminum-Oxygen cell configuration.
- Figure 7. Absorption spectra of a 1.05:1 $\text{AlCl}_3\text{:BuPyCl}$ (dashed line) and 0.96 mM $\text{Ru}(\text{bpy})_3^{2+}$ in 1.05:1 $\text{AlCl}_3\text{:BuPyCl}$ melt (solid line). Cell thickness = 0.1 cm.
- Figure 8. Absorption spectra of: 1.17 mM $\text{Fe}(\text{bpy})_3^{2+}$ in 1:1 melt, cell thickness = 0.1 cm (---); 2.5 mM $\text{Fe}(\text{phen})_3^{3+}$ in 1.9:1 melt, cell thickness = 1 cm (—).

Figure 9. Absorption spectra of: 1.45 mM $\text{Fe}(\text{phen})_3^{2+}$ in 1:1 melt, cell thickness = 0.1 cm (---); 1.7 mM $\text{Ru}(\text{bpy})_3^{3+}$ in 1.9:1 melt, cell thickness = 0.1 cm (-.-); 3.87 mM $\text{Fe}(\text{bpy})_3^{3+}$ in 1.2:1 melt, cell thickness = 1 cm (—).




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9

DL/413/83/01
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Dr. M. B. Denton
Department of Chemistry
University of Arizona
Tucson, Arizona 85721

Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. J. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. B. R. Kowalski
Department of Chemistry
University of Washington
Seattle, Washington 98105

Dr. H. Freiser
Department of Chemistry
University of Arizona
Tucson, Arizona 85721

Dr. H. Chernoff
Department of Mathematics
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. A. Zirino
Naval Undersea Center
San Diego, California 92132

Professor George H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. Alan Bewick
Department of Chemistry
Southampton University
Southampton, Hampshire
ENGLAND 5095NA

Dr. S. P. Perone
Lawrence Livermore Laboratory L-370
P.O. Box 808
Livermore, California 94550

Dr. L. Jarvis
Code 6100
Naval Research Laboratory
Washington, D.C. 20375

Dr. G. M. Hieftje
Department of Chemistry
Indiana University
Bloomington, Indiana 47401

Dr. Christie G. Enke
Department of Chemistry
Michigan State University
East Lansing, Michigan 48824

Dr. D. L. Venezky
Naval Research Laboratory
Code 6130
Washington, D.C. 20375

Walter G. Cox, Code 3632
Naval Underwater Systems Center
Building 148
Newport, Rhode Island 02840

Professor Isiah M. Warner
Department of Chemistry
Emory University
Atlanta, Georgia 30322

Dr. Kent Eisentraut
Air Force Materials Laboratory
Wright-Patterson AFB, Ohio 45433

Dr. Adolph B. Amster
Chemistry Division
Naval Weapons Center
China Lake, California 93555

Dr. B. E. Oouda
Chemical Sciences Branch
Code 50 C
Naval Weapons Support Center
Crane, Indiana 47322

Dr. John Eyler
Department of Chemistry
University of Florida
Gainesville, Florida 32611

DL/413/83/01
051B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Professor J. Janata
Department of Bioengineering
University of Utah
Salt Lake City, Utah 84112

Dr. J. DeCorpo
NAVSEA
Code 05R14
Washington, D.C. 20362

Dr. Charles Anderson
Analytical Chemistry Division
Athens Environmental Laboratory
College Station Road
Athens, Georgia 30613

Dr. Ron Flemming
B 108 Reactor
National Bureau of Standards
Washington, D.C. 20234

Dr. David M. Hercules
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania

Dr. Frank Herr
Office of Naval Research
Code 422CB
800 N. Quincy Street
Arlington, Virginia 22217

Professor E. Keating
Department of Mechanical Engineering
U.S. Naval Academy
Annapolis, Maryland 21401

Dr. M. H. Miller
1133 Hampton Road
Route 4
U.S. Naval Academy
Annapolis, Maryland 21401

Dr. Clifford Spiegelman
National Bureau of Standards
Room A337 Bldg. 101
Washington, D.C. 20234

Dr. Denton Elliott
AFOSR/NC
Bolling AFB
Washington, D.C. 20362

Dr. B. E. Spielvogel
Inorganic and Analytical Branch
P.O. Box 12211
Research Triangle Park, NC 27709

Ms. Ann De Witt
Material Science Department
160 Fieldcrest Avenue
Raritan Center
Edison, New Jersey 08818

Dr. A. Harvey
Code 6110
Naval Research Laboratory
Washington, D.C. 20375

Dr. John Hoffsommer
Naval Surface Weapons Center
Building 30 Room 208
Silver Spring, Maryland 20910

Mr. S. M. Hurley
Naval Facilities Engineering Command
Code 032P
200 Stovall Street
Alexandria, Virginia 22331

Ms. W. Parkhurst
Naval Surface Weapons Center
Code R33
Silver Spring, Maryland 20910

Dr. M. Robertson
Electrochemical Power Sources Division
Code 305
Naval Weapons Support Center
Crane, Indiana 47522

CDR Andrew T. Zander
10 Country Club Lane
ONR Boston
Plaistow, New Hampshire 03865

DL/413/83/01
051B/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 051B

Dr. Robert W. Shaw
U.S. Army Research Office
Box 12211
Research Triangle Park, NC 27709

Dean William Tolles
Naval Post Graduate School
Spanaugel Hall
Monterey, California 93940

Dr. Marvin Wilkerson
Naval Weapons Support Center
Code 30511
Crane, Indiana 47522

Dr. H. Wohltjen
Naval Research Laboratory
Code 6170
Washington, D.C. 20375

Dr. J. Wyatt
Naval Research Laboratory
Code 6110
Washington, D.C. 20375

DL/413/83/01
359/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Paul Delahay
Department of Chemistry
New York University
New York, New York 10003

Dr. P. J. Hendra
Department of Chemistry
University of Southampton
Southampton SO9 5NH
United Kingdom

Dr. T. Katan
Lockheed Missiles and
Space Co., Inc.
P.O. Box 504
Sunnyvale, California 94088

Dr. D. N. Bennion
Department of Chemical Engineering
Brighma Young University
Provo, Utah 84602

Dr. R. A. Marcus
Department of Chemistry
California Institute of Technology
Pasadena, California 91125

Mr. Joseph McCartney
Code 7121
Naval Ocean Systems Center
San Diego, California 92152

Dr. J. J. Auburn
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Joseph Singer, Code 302-1
NASA-Lewis
21000 Brookpark Road
Cleveland, Ohio 44135

Dr. P. P. Schmidt
Department of Chemistry
Oakland University
Rochester, Michigan 48063

Dr. H. Richtol
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. C. E. Mueller
The Electrochemistry Branch
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, Maryland 20910

Dr. Sam Perone
Chemistry & Materials
Science Department
Lawrence Livermore National Lab.
Livermore, California 94550

Dr. Royce W. Murray
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. G. Goodman
Johnson Controls
5757 North Green Bay Avenue
Milwaukee, Wisconsin 53201

Dr. B. Brummer
EIC Incorporated
111 Chapel Street
Newton, Massachusetts 02158

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Electrochimica Corporation
Attn: Technical Library
2485 Charleston Road
Mountain View, California 94040

Library
Duracell, Inc.
Burlington, Massachusetts 01803

Dr. A. B. Ellis
Chemistry Department
University of Wisconsin
Madison, Wisconsin 53706

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. M. Wrighton
Chemistry Department
Massachusetts Institute
of Technology
Cambridge, Massachusetts 02139

Dr. B. Stanley Pons
Department of Chemistry
University of Utah
Salt Lake City, Utah 84112

Donald E. Mains
Naval Weapons Support Center
Electrochemical Power Sources Division
Crane, Indiana 47522

S. Ruby
DOE (STOR)
M.S. 68025 Forrestal Bldg.
Washington, D.C. 20595

Dr. A. J. Bard
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. Janet Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214

Dr. Donald W. Ernst
Naval Surface Weapons Center
Code R-33
White Oak Laboratory
Silver Spring, Maryland 20910

Mr. James R. Moden
Naval Underwater Systems Center
Code 3632
Newport, Rhode Island 02840

Dr. Bernard Spielvogel
U.S. Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

Dr. William Ayers
ECD Inc.
P.O. Box 5357
North Branch, New Jersey 08876

Dr. M. M. Nicholson
Electronics Research Center
Rockwell International
3370 Miraloma Avenue
Anaheim, California

Dr. Michael J. Weaver
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. R. David Rauh
EIC Corporation
111 Chapel Street
Newton, Massachusetts 02158

Dr. Aaron Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH ENGLAND

~~Dr. R. A. Osteryoung
Department of Chemistry
State University of New York
Buffalo, New York 14214~~

Dr. Denton Elliott
Air Force Office of Scientific
Research
Bolling AFB
Washington, D.C. 20332

Dr. R. Nowak
Naval Research Laboratory
Code 6130
Washington, D.C. 20375

Dr. D. F. Shriver
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Aaron Fletcher
Naval Weapons Center
Code 3852
China Lake, California 93555

DL/413/83/01
359/413-2

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. David Aikens
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. A. P. B. Lever
Chemistry Department
York University
Downsview, Ontario M3J1P3

Dr. Stanislaw Szpak
Naval Ocean Systems Center
Loc: 6343, Bayside
San Diego, California 95152

Dr. Gregory Farrington
Department of Materials Science
and Engineering
University of Pennsylvania
Philadelphia, Pennsylvania 19104

M. L. Robertson
Manager, Electrochemical
and Power Sources Division
Naval Weapons Support Center
Crane, Indiana 47522

Dr. T. Marks
Department of Chemistry
Northwestern University
Evanston, Illinois 60201

Dr. Micha Tomkiewicz
Department of Physics
Brooklyn College
Brooklyn, New York 11210

Dr. Lesser Blum
Department of Physics
University of Puerto Rico
Rio Piedras, Puerto Rico 00931

Dr. Joseph Gordon, II
IBM Corporation
K33/281
5600 Cottle Road
San Jose, California 95193

Dr. D. H. Whitmore
Department of Materials Science
Northwestern University
Evanston, Illinois 60201

Dr. Alan Bewick
Department of Chemistry
The University of Southampton
Southampton, SO9 5NH ENGLAND

Dr. E. Anderson
NAVSEA-56233 NC #4
2541 Jefferson Davis Highway
Arlington, Virginia 20362

Dr. Bruce Dunn
Department of Engineering &
Applied Science
University of California
Los Angeles, California 90024

Dr. Elton Cairns
Energy & Environment Division
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

Dr. D. Cipris
Allied Corporation
P.O. Box 3000R
Morristown, New Jersey 07960

Dr. M. Philpott
IBM Corporation
5600 Cottle Road
San Jose, California 95193

Dr. Donald Sandstrom
Department of Physics
Washington State University
Pullman, Washington 99164

Dr. Carl Kannewurf
Department of Electrical Engineering
and Computer Science
Northwestern University
Evanston, Illinois 60201

TECHNICAL REPORT DISTRIBUTION LIST, 359

Dr. Robert Somoano
Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California 91103

Dr. Johann A. Joebstl
USA Mobility Equipment R&D Command
DRDME-EC
Fort Belvoir, Virginia 22060

Dr. Judith H. Ambrus
NASA Headquarters
M.S. RTS-6
Washington, D.C. 20546

Dr. Albert R. Landgrebe
U.S. Department of Energy
M.S. 6B025 Forrestal Building
Washington, D.C. 20595

Dr. J. J. Brophy
Department of Physics
University of Utah
Salt Lake City, Utah 84112

Dr. Charles Martin
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Dr. H. Tachikawa
Department of Chemistry
Jackson State University
Jackson, Mississippi 39217

Dr. Theodore Beck
Electrochemical Technology Corp.
3935 Leary Way N.W.
Seattle, Washington 98107

Dr. Farrell Lytle
Boeing Engineering and
Construction Engineers
P.O. Box 3707
Seattle, Washington 98124

Dr. Robert Gotscholl
U.S. Department of Energy
MS G-226
Washington, D.C. 20545

Dr. Edward Fletcher
Department of Mechanical Engineering
University of Minnesota
Minneapolis, Minnesota 55455

Dr. John Fontanella
Department of Physics
U.S. Naval Academy
Annapolis, Maryland 21402

Dr. Martha Greenblatt
Department of Chemistry
Rutgers University
New Brunswick, New Jersey 08903

Dr. John Wasson
Syntheco, Inc.
Rte 6 - Industrial Pike Road
Gastonia, North Carolina 28052

Dr. Walter Roth
Department of Physics
State University of New York
Albany, New York 12222

Dr. Anthony Sammells
Eltron Research Inc.
710 E. Ogden Avenue #108
Naperville, Illinois 60540

Dr. W. M. Risen
Department of Chemistry
Brown University
Providence, Rhode Island 02192

Dr. C. A. Angell
Department of Chemistry
Purdue University
West Lafayette, Indiana 47907

Dr. Thomas Davis
Polymer Science and Standards
Division
National Bureau of Standards
Washington, D.C. 20234

ATE
LME